
1996 Annual Report, submitted electronically

Heidi M. Sosik and Robert J. Olson

Response of Particulate Optical Properties to Coastal Mixing Processes

Biology Department, MS 32 Woods Hole Oceanographic Institution Woods Hole, MA 02543-1049 hsosik@whoi.edu 508-289-2311 508-457-2134



Long-Term Goals

Our long-term goals are to develop a better understanding of the relationships between upper ocean optical properties and particulate and dissolved seawater constituents, and to determine how these relationships are influenced by physical processes. Specific goals include both predicting and modeling optical variability relevant for biological processes, such as phytoplankton photosynthesis, and retrieval of information about the biomass and activity of plankton from optical measurements.

Scientific/Technical Objectives

Spatial and temporal variability in particulate and dissolved material is a significant source of optical variability in the upper ocean. The primary objective of the present work is to examine the interaction between physical processes and the properties, abundance, and optical significance of different particle types in coastal ocean waters. Specific project objectives are to refine individual particle measurement methods and develop approaches to using individual particle results for interpretation of both inherent and apparent bulk optical properties. The project includes a combination of instrument development and field studies in coastal waters of the eastern US continental shelf.

Approach

The approach is to use techniques for characterizing and assessing the optical properties of particles using both *in situ* and ship-board instrumentation and both bulk and single particle methods. Our primary tools are flow cytometry for individual particle light scattering and fluorescence properties, spectrophotometry for measuring bulk dissolved and particulate absorption spectra (including separation of phytoplankton pigment absorption from the bulk based on methanol extraction), and spectral underwater radiometry. The flow cytometric and spectrophotometric measurements will be conducted both on discrete water samples and in situ. In situ measurement provides the opportunity for relatively unperturbed sampling, with generally greater spatial resolution, while analysis of discrete water samples continues to provide more detailed characterization of optically-active seawater constituents.

Tasks Completed

 We have completed modifications to our existing EPICS flow cytometer, expanding the dynamic range for forward and side angle scattering and chlorophyll

19970324 028

fluorescence measurements by optically dividing and directing the signals to separate photomultipliers with different sensitivities. These modifications are aimed at streamlining sample analysis which previously required separate instrument configurations to cover the particle range of < 1 to ~ 20 microns in diameter. We have also been continuing development of our in situ flow cytometer (independently funded by NSF and DOE), aiming for deployment during the Spring 1997 field program.

2) A large and interesting dataset was collected on the first cruise of this project during August - September 1996. We conducted roughly 3 vertical profiles daily for 3 weeks at the Coastal Mixing and Optics central experiment site on the continental shelf south of Woods Hole. Discrete water sample analysis included complete flow cytometric characterization of the phytoplankton and other particles between approximately 0.7 and 20 microns in diameter. Vertical profiles of spectral downwelling irradiance and upwelling radiance were collected with a free-fall instrument (Satlantic) deployed away from the ship. All instruments performed very well and data analysis is currently underway.

Results

Preliminary analysis of results from the late summer cruise show unexpectedly high levels of day-to-day variability throughout the sampling period, which peaked with the passage of hurricane Eduoard through the study site. The decrease in stratification following the hurricane was accompanied by changes in the optical properties of the water column (Fig. 1), attributable primarily to changes in the abundance and characteristics of suspended material.

Flow cytometric analysis of particle properties (Fig. 2) on the day before the hurricane indicated that at the surface, phytoplankton and non-fluorescent particles accounted for nearly equal amounts of forward light scattering. Light scattering in the chlorophyll maximum layer (27 m) was dominated by phytoplankton, and near the bottom, phytoplankton were much less important and non-fluorescent particles much more, as expected.

During the hurricane the water column may have mixed completely, although two days afterwards, when we sampled the site again, both hydrography and particle distributions were showing evidence of re-stratification. The size distributions of light scattering by phytoplankton at the surface and at 20 m, for example, were dramatically different, with a large increase in the ultraphytoplankton (2-5 μ m cells) at the surface. In addition, the size distributions were markedly different from those of the previous sampling: Cells at the large end of the size distribution (>10 μ m) had become relatively less abundant, and at the surface the smallest cells (*Synechococcus*) had increased in size.

After the hurricane, non-fluorescent particles had become the major contributors to light scattering in the chlorophyll maximum layer as well as near the bottom; at the surface there was almost no change. In the subsurface samples, the size distribution of the non-fluorescent particles was shifted dramatically to smaller sizes, from a mode of about 10 μm to about 2 μm .

This is an example of the type of analysis which we will be pursuing for our entire data set, spanning a 3 week time period and including several profiles each day.

Impacts for Science and Technology and/or Applications

This project includes the development of improved techniques for analyzing marine particles and characterizing their optical properties. Our ability to independently quantify size distributions for phytoplankton and non-phytoplankton particles is a new contribution which will lead to better understanding of optical variability in the ocean.

Relationships to Other Projects

This project is closely tied to Sosik's NASA New Investigator Program award. The NASA support will be used to investigate the regulation of local biological production of particles at the CMO site and to explore the effects of changes in particle properties on ocean color. In addition, Olson is independently funded (DOE, NSF) to develop an in situ flow cytometer which will be deployed at the CMO site on future cruises. The interpretation of flow cytometric light scattering and fluorescence distributions is also being supported by Olson's NSF JGOFS project for work in the Arabian Sea.




Figure 1. Hydrographic and optical data collected on the first CMO Optics cruise at stations near the central mooring site (\sim 65 m water depth), before (August 31) and after (September 4) the passage of hurricane Eduoard. Vertical distributions of temperature and salinity were relatively uniform after the hurricane, but variations in diffuse attenuation were high. The subsurface peak in K_d at 490 nm on August 31 is associated with phytoplankton, as indicated by the pronounced minimum in K_d at 683 nm (due to chlorophyll fluorescence), while the high attenuation near the bottom on September 4 is due to resuspended material from the bottom.




Figure 2. Particle size distributions for phytoplankton and non-phytoplankton showing relative contributions to forward light scattering (FLS), as measured by the flow cytometer. Water was collected from discrete depths for the same 2 days as in Figure 1 and approximately 5 ml samples were analyzed. FLS values are expressed in comparable arbitrary units in all cases. Note the change in scale for the non-phytoplankton particles.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average I hour per response, including the time for reviewing instructions, learning existing data sources, gathering and maintaining the data needed, and commetting and reviewing the collection of information. Send commettin regarding this burden estimate or any other aspect of this collection of information, including suppersions for reducing this burden. In Washington Heaviers Services, Directorate for Information Collection and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA. 22202.4302, and to the Office of Management and Budget, Paperwork Reduction Project (1704-0188), Washington, OC. 2003.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	PROGRESS RE	DATES C	OVERED 2/1/96-
A TITLE AND SUBTITLE				NG NUMBERS
RESPONSE OF PARTICULATE OPTICAL PROPERTIES TO COASTAL MIXING PROCESSES			N00014-95-1-0333	
6. AUTHOR(S)				
Heidi Sosik and Robert J. Olson				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFO	RMING ORGANIZATION
WOODS HOLE OCEANOGRAPHIC INSTITUTION				T NUMBER
WOODS HOLE, MA 02543			WHO)I NO. 9364
·			.,,,,,	2 7504
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10 25 5	
Dr. Steven Ackleson				SORING/MONITORING CY REPORT NUMBER
Code 322, ONR				
Ballston Tower One				
800 N. Quincy St, Arlington, VA 22217-5660				
11. SUPPLEMENTARY NOTES				
IL SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STA	TEMENT		12b. DIST	RIBUTION CODE
Approved for Public Release; Distribution is				
Unlimited.				
on a time cod .			}	
13. ABSTRACT (Maximum 200 words)			L	
Spatial and temporal variability in particulate and dissolved material is a significant source of				
optical variability in the upper ocean. The goal of this project is to examine the interaction between				
physical processes and the properties, abundance, and optical significance of different particle types in				
coastal ocean waters. Specific project objectives are to refine individual particle measurement methods				
and develop approaches to use individual particle results for interpretation of both inherent and apparent				
bulk optical properties. The project includes a combination of instrument development and field studies in				
waters of the eastern US continental shelf as part of the Coastal Mixing and Optics Accelerated Research				
Initiative. Techniques including bulk absorption measurements and individual particle analysis are being				
used and sampling includes a combination of in situ measurements and ship-board analysis of water.				
Absorption is evaluated as a function of categories distinguished based on filtration (particulate vs.				
dissolved) and by chemical methods (phytoplankton vs. detritus). Distributions of particle size and scatter-				
ing are determined for various particle types (including phytoplankton and non-phytoplankton) which can				
be distinguished by their flow cytometric fluorescence and scattering properties. A 10-month experiment				
is currently underway, with one 3-week cruise completed and a second planned for April-May, 1997.				
14. SUBJECT TERMS				15. NUMBER OF PAGES
				16. PRICE CODE
17. SECURITY CLASSIFICATION 118.	SECURITY CLASSIFICATION	19 SECTION CLASS	CATION	20 111474 7100 00 1007
· OF REPORT	OF THIS PAGE	19. SECURITY CLASSIF OF ABSTRACT	CATION	20. LIMITATION OF ABSTRACT
NSN 7540-01-280-5500 Standard Form 298 (Rev. 2-89)				
•			29	escribed by ANSI Std. 239-18 8-102